# Lecture 21 Impacts and Extinction



Astronomy 141 - Autumn 2012

This lecture examines the role of asteroidal impacts on the history of life on Earth.



Major impacts ended 3.8 Gya at the end of the Hadean Eon, but impacts generally tapered off to a steady rate.



Near Earth Asteroids (NEOs) are asteroids that orbit near the Earth, with 1000-2000 ~1 km in size.



An asteroidal impact caused the extinction of the dinosaurs 65Mya, and perhaps other mass die offs.




5 major mass extinctions in the past 500 Myr important for driving diversity and evolution.

Impacts by 300-500 km asteroids during the Hadean would have sterilized the early Earth.


Impact energy would have vaporized the oceans and melted much of the crust.

"Steam atmospheres and magma seas."

Any early life would have been wiped out.



The last such sterilizing impacts occurred during the Hadean Eon  $4.3-3.8\;\mbox{Gya}$ 



Craters from past impacts are found on all continents, with 50 craters larger than 20km across.

Barringer Crater Arizona
50m asteroid, 50kya

Largest is Vredefort Crater
S. Africa, 10km asteroid
that struck 2.2 Gya


Lake Manicouagan
5km asteroid 214 ± 1Mya

The sources of asteroids today are the asteroid belt and outer solar system.

Asteroids are leftover rocky/icy material from the formation of the Solar System

Disturbed by Jupiter's gravity into the inner solar system.

~1.2 million asteroids larger than 1km across, but only 100 more than 140 km across.



Near-Earth Objects (NEOs) are asteroids and comets that have orbits that bring them near Earth. 6300 known, most a few meters across. Probably 1000-2000 NEOs larger than 1 km across. On unstable orbits, but are constantly supplied from the outer asteroid belt. Numerous groups are actively searching for NEOs. (Spaceguard) A hit by the small asteroid 2008 TC3 was predicted and occurred in the Sudan in October 2008. First prediction of a small asteroid strike - but only 20hr before impact. Asteroid was 3-5m in size. Burned up in the atmosphere, but 4 kg of fragments were collected. Energy release was 1-2 kilotons. Rocks of this size hit the Earth 2 or 3 times per year. A bigger direct hit occurred in Siberia in 1908 near the Tunguska River Seismic activity recorded 1000 km away. Airburst ~10-15 km altitude with an estimated energy release of 10-15 megatons. Probably an asteroid or comet 40 meters in diameter. Such strikes should happen

once per century.

The K-T Boundary is a layer of sediments that divides the Cretaceous and Tertiary Periods Boundary occurs 65 Myr ago. Layer of clay sediment unusually rich in Iridium and soot. The Iridium content is 100x the natural abundance in the crust, but comparable to what is found in asteroids and chondrites. At the time of the K-T boundary 65 Mya, 75% of land and marine species, including all non-avian dinosaurs, went extinct. The favored explanation is that the K-T boundary is the after effects of a massive asteroidal impact. 10 km diameter asteroid making a direct hit. Energy release 2 million times that of the most powerful nuclear bombs. Global firestorms Mega tsunamis Dust in the air causing 10-20% less sunlight. Physical evidence of the impact is Chicxulub Crater in the Yucatan. 180 km in diameter. Shocked quartz Impact glasses (tektites) Gravitational anomaly Radiometric dating

gives 65 Mya

